

PHARMACOPŒIA




GUY'S HOSPITAL

M

8917

22500288097




Sister Charity.

April 1st 1886.

From S S R.

R. M. Charles

Guys Hospital.



FORMULÆ

USED AT

GUY'S HOSPITAL,

IN ADDITION TO THOSE IN THE

BRITISH PHARMACOPŒIA.

COMPILED BY

A COMMITTEE OF THE STAFF.

LONDON:


J. & A. CHURCHILL, NEW BURLINGTON STREET.

1879.

24 876 043

102-4

WELLCOME INSTITUTE LIBRARY	
Coll.	<u>Wellcome</u>
Coll.	<u>Gen. Coll.</u>
No.	<u>QV740</u>
	<u>M</u>
	<u>89171</u>
	<u>G</u>



GUY'S HOSPITAL.

MEDICAL AND SURGICAL STAFF.

Consulting Physicians.

Sir William Gull, Bart., M.D., D.C.L., F.R.S.;

G. Owen Rees, M.D., F.R.S.

Physicians.

S. O. Habershon, M.D.; S. Wilks, M.D., F.R.S.;

F. W. Pavy, M.D., F.R.S.; W. Moxon, M.D.

Assistant Physicians.

C. Hilton Fagge, M.D.; P. H. Pye-Smith, M.D.;

Frederick Taylor, M.D.; J. F. Goodhart, M.D.

Consulting Surgeons.

E. Cock, F.R.C.S.; J. Birkett, F.R.C.S.

Surgeons.

J. Cooper Forster, F.R.C.S.; Thomas Bryant, F.R.C.S.;

Arthur Durham, F.R.C.S.; H. G. Howse, M.S.

Assistant Surgeons.

N. Davies-Colley, M.C.; R. Clement Lucas, B.S.;

C. H. Golding-Bird, M.B.; W. H. A. Jacobson, M.B.

Consulting Obstetric Physician.—Henry Oldham, M.D.

Obstetric Physician.—J. Braxton Hicks, M.D., F.R.S.

Assistant Obstetric Physician.—A. L. Galabin, M.D.


Ophthalmic Surgeon.—C. Bader, Esq.

Assistant Ophthalmic Surgeon.—C. Higgins, F.R.C.S.

Dental Surgeon.—S. J. A. Salter, M.B., F.R.S.

Assistant Dental Surgeon.—H. Moon, Esq.

Aural Surgeon.—W. Laidlaw Purves, M.D.



Ammon. Brom. gr xxv
Chlorae Hydr. gr xxv
Aqua ai 3i. o. n.

Sig. M. Peruviana.
1/2 the strength of Condys Fluid
gr 4 to 3i, dilute in 40 parts
water for gargle

For ear-ache

J. Pinum & Glycerin a-a. hot

For *Chelblain { also
*Zinc Oxide 3i
Glycerin 3ss
Gaulther 3i 3iii

Apply equal parts of

J. Opie

Spirits of Camphor -

Bloway, for Phleto. (Horse)

Blow. Wax lb i
Ol Carbol 3ss

Heat but do not boil -

GUY'S HOSPITAL

FORMULÆ.



AQUA GLYCYRRHIZÆ.

Liquorice Water.

Extract of Liquorice	.	.	.	$\frac{1}{2}$ oz.
Water	.	.	.	1 pint

Dissolve.

ARGENTI NITRAS MITIOR.

Milder Nitrate of Silver.

Nitrate of Silver	.	.	.	1 part
Nitrate of Potash	.	.	.	3 parts

Fuse together and cast in moulds.
For local application.

BALNEUM AËRIS CALIDI.

Warm Air Bath.

120° to 150° Fahr.

BALNEUM AQUÆ FRIGIDÆ.

Cold Water Bath.

60° Fahr.

BALNEUM AQUÆ TEPIDÆ.

Tepid Water Bath.

85° to 92° Fahr.

BALNEUM AQUÆ CALIDÆ.

Warm Water Bath.

98° to 110° Fahr.

BALNEUM ALKALINUM.

Alkaline Bath.

Carbonate of Soda . . . 4 oz.

Warm Water . . . 30 gallons

Dissolve.

BALNEUM ACIDI NITRO-HYDRO-
CHLORICI.*Nitro-Hydrochloric Acid Bath.*

Nitric Acid . . . 1 lb.

Hydrochloric Acid . . . 1½ lbs.

Warm Water . . . 30 gallons

Mix.


Imperial Drink

Grain of Tartar 31- if

Ag ad 67 boiling water

add lemon juice & Sugar to flavor.

(if allowed)



CATAPLASMA CAROTÆ.

Carrot Poultice.

Common Carrots	}	. . .	a sufficiency.
Water			

Boil the Carrots until soft, then pulp them.

CODEIA.

An Alkaloid obtained from Opium.

Dose.— $\frac{1}{2}$ to 5 grs.

COLLYRIUM ALUMINIS.

Alum Eye Wash.

Alum		3 grs.
Distilled Water		1 fl. oz.

Dissolve.

COLLYRIUM BELLADONNÆ.

Belladonna Eye Wash.

Belladonna Extract		8 grs.
Distilled Water		1 fl. oz.

Mix.

COLLYRIUM ZINCI CHLORIDI.

Chloride of Zinc Eye Wash.

Chloride of Zinc		2 grs.
Distilled Water		1 fl. oz.

Dissolve.

COLLYRIUM ZINCI SULPHATIS.

Sulphate of Zinc Eye Wash.

The same as the Sulphate of Zinc Lotion.
(2 grs. to 1 fl. oz.)

CONFECTIO SENNÆ ET SULPHURIS.

Confection of Senna and Sulphur.

Confection of Senna	}	.	.	equal parts.
Confection of Sulphur				

Mix thoroughly.

Dose.—1 to 2 *drs.*

COPAIBÆ RESINA.

Resin of Copaiva.

Made by heating Copaiba over a water bath until the oil is dissipated and a hard resin remains.

When pure, the resin is of an amber colour : if green, copper is present as an impurity.

Used in the Resin of Copaiva Mixture.

CURARE (WOORALI).

The inspissated juice of the Strychnos Toxicifera.


Used in the Injection of Curare.

DATURINA.

Daturine.

An alkaloid obtained from the Datura Stramonium.

Used in the Ointment of Daturine.



ELECTRICITAS.

1. FRICTIONAL, STATIC, or FRANKLINIC
ELECTRICITY.

The electricity in small quantity, but of high tension, which is produced by friction from the rotatory glass disc of an electrical machine.

The patient is placed on an insulated stool, and is charged from the machine by a metallic conductor. Sparks are then taken with a metallic conductor from the part of the body directed.

Static electricity may also be applied as an electric shock from a Leyden jar, to be discharged as directed by means of insulated conductors.

2. GALVANISM, or VOLTAIC ELECTRICITY.

The current produced by the chemical action of a galvanic battery, the elements of which are arranged in series.


The electromotive force will depend on the number of elements used. In the Leclanché battery, now used in the Hospital, from five to thirty are employed. In the Muirhead battery ten to forty are usually employed.

The Galvanic Current is applied by means

of insulated conducting wires and electrodes (poles or rheophores). These may be metal plates, plates covered with leather, moist sponges in holders, wire brushes, &c.; moist surfaces being used for bringing the current to deeper parts, and dry points for affecting the skin.

Galvanism is applied in two ways—as a continuous and as an interrupted current. The “constancy” of the current means in both cases its freedom from minor variations of intensity and depends on the construction and working of the battery used.

(a) When *Continuous Galvanism* is ordered, the two electrodes, of leather or sponge moistened with salt and water, must be kept applied to the parts directed. The positive electrode or anode (connected with the positive pole of the battery) is usually placed on the spine or the proximal part of the muscle or nerve, the negative electrode or cathode (connected with the negative pole of the battery) on the distal part, for a period of five or ten minutes or longer. The skin should be well sponged with salt and water, so as to render it as good a conductor as possible. Or the negative electrode may be dipped in a tepid salt-water foot or hand bath, the water serving as the conductor.



(b) When *Interrupted Galvanism* is ordered, one electrode (usually the anode, but this should be directed according as the ascending or descending current is desired) is kept applied to the spine or over some proximal point of the muscle or nerve to be acted on, while the other is moved over the muscle or nerve, either quickly (once every second) or more slowly. Or, both electrodes being held fixed on the parts directed, the circuit may be made and broken at another point.

3. FARADISM, or INDUCED ELECTRICITY.

The *secondary* or induced currents evolved from an induction coil connected with a battery of few elements. This current is of considerable electromotive force. The *primary* or galvanic current from the battery to the induction coil is made and broken by a magnetic interrupter. The induced current is "alternating," that is, alternately ascending and descending. It is not electrolytic.

To be applied by means of suitable electrodes kept in contact with the parts directed, or slowly moved over them.

4. THE GALVANIC CAUTERY.

A platinum wire raised to a dull red heat by

means of a galvanic current of considerable strength, obtained from a Bunsen's battery of a few large elements arranged as one.

The current is passed through two copper electrodes to a loop of platinum wire included in the circuit. This wire loop may be very narrow (platinum point), or be wound round a small piece of porcelain (porcelain button), or be used in the form of an *écraseur* (galvanic *écraseur*).

ENEMA COLOCYNTHIDIS COMPOSITUM.

Compound Colocynth Enema.

Colocynth Pulp, cut	60 grs.
Water	12 fl. oz.

Boil for ten minutes and strain.

Then add

Common Salt	$\frac{1}{2}$ oz.
-----------------------	-------------------


And mix.

ENEMA COMMUNE.

Common Enema.

Common Salt	1 oz.
Oatmeal Gruel	12 fl. oz.

Dissolve.



Emema opii.

Tinct. opii 3p-

Starch 3*ij*.

Mucilage of Starch.

Starch 1-

Distilled water 40-

Boil & stirring, for a few minutes.

Emema Terebinthinae

Oil of Turpentine 3*i*.

Mucilage of Starch 3*xv*

Emema Mag. Sulph

Sulphate of Magnesia in Crystals 3*i*
Dissolve in 3*xv* of Mucilage of Starch
add 3*i* of Olive oil.

ENEMA CIBI.

Nutrient Enema.

Warm Beef-tea 2 fl. oz.
One egg.

Mix thoroughly.

Brandy or Opium to be added as required.

ENEMA OLEI.

Olive Oil Enema.

Olive Oil 4 fl. oz.
Warm Mucilage of Starch 8 fl. oz.

Mix thoroughly.

ENEMA OLEI RICINI.

Castor Oil Enema.

Castor Oil 1 fl. oz.
Warm Mucilage of Starch 11 fl. oz.

Mix thoroughly.

ENEMA SAPONIS.

Soap Enema.

Soft Soap 1 oz.
Boiling Water 16 fl. oz.

Dissolve, and use when sufficiently cool.

ERGOTINA.

Ergotine (Bonjean's).

*The active matter of Ergot of Rye in a
concentrated form.*

Used in the Injection of Ergotine.

ESERINE (OR PHYSOSTIGMINÈ).

The active principle of Calabar Bean.

The Sulphate of Eserine as used in the Ointment and Solution of Eserine.

FERRI ET AMMONIÆ TARTRAS.

Ammonio-Tartrate of Iron.

Dose.—5 to 10 grs.

GARGARISMA ACIDI CARBOLICI.

Carbolic Acid Gargle.

Carbolic Acid (No. 2 Calvert)	.	3 grs.
Distilled Water	.	1 fl. oz.

Dissolve.

GARGARISMA ACIDI HYDROCHLORICI.

Hydrochloric Acid Gargle.

The same as the Mixture. (10 m. to 1 fl. oz.)

GARGARISMA ACIDI TANNICI.

Tannic Acid Gargle.

The same as the Lotion. (10 grs. to 1 fl. oz.)

GARGARISMA ALUMINIS.

Alum Gargle.


Alum	.	.	.	.	.	5 grs.
Water	.	.	.	.	.	1 fl. oz.

Dissolve.

Acid Sulphuricos 3℥
 Glycerine 3℥
 Aqua ad 3℥ - pro: garg.

Mouth wash (Mr. Durham)

Myrrh	1 Part
Ess. de Clove	16 "
Borax	1 "
Water	3 "
Syrup	3 "



GARGARISMA ALUMINIS COMPOSITUM.

Compound Alum Gargle.

Alum	5 grs.
Syrup of Red Roses	1 fl. dr.
Decoction of Yellow Cinchona Bark	7 fl. drs.

Dissolve the alum in the decoction, then add the syrup.

GARGARISMA BORACIS.

Boræ Gargle.

Borax, powdered	5 grs.
Syrup of Red Roses	1 fl. dr.
Barley Water	7 fl. drs.

Mix.

GARGARISMA POTASSÆ CHLORATIS.

Chlorate of Potash Gargle.

The same as the Mixture. (10 grs. to 1 fl. oz.)

GELATINUM ATROPIÆ.

Atropine Gelatine.

For local application in the form of discs.

GELATINUM ESERINÆ.

Eserine Gelatine.

For local application.

GELATINUM PHYSOSTIGMATIS FABÆ.

Calabar Bean Gelatine.

For local application.

GLYCERINUM PLUMBI ACETATIS.

Glycerine of Acetate of Lead.

Acetate of Lead 1 dr.

Glycerine 1 fl. oz.

Dissolve.

For local application.

GUTTÆ ARGENTI NITRATIS.

Nitrate of Silver Drops.

Nitrate of Silver 1 gr.

Distilled Water 1 fl. oz.

Dissolve.

GUTTÆ ATROPIÆ.

Atropine Drops.

Sulphate of Atropine $\frac{1}{2}$ gr.

Distilled Water 1 fl. oz.

Dissolve.


GUTTÆ CUPRI SULPHATIS.

Sulphate of Copper Drops.

Sulphate of Copper 2 grs.

Distilled Water 1 fl. oz.

Dissolve.



GUTTÆ OPII.

Opium Drops.

Wine of Opium 2 fl. drs.

Distilled Water 3 fl. drs.

Mix.

HYDRARGYRI OLEAS.

Oleate of Mercury.

This contains 10 per cent. of Oxide of Mercury.

For local application.

HYDRARGYRI OLEAS CUM MORPHIA.

Oleate of Mercury with Morphia.

Morphia 1 gr.

Oleate of Mercury 1 dr.

Dissolve.

For local application.

HYDRARGYRI SUBOXIDUM.

(Syn. Hydrargyri Oxidum Nigrum vel Cinereum.)

Grey or Mercurous Oxide.

Dose.—1 to 5 grs.

INHALATIO.

The Inhalation of a Medicated Steam Spray.

The following are of a strength to be used with a Seigel's Steam Spray Producer:—

INHALATIO ACIDI CARBOLICI.

Inhalation of Carbolic Acid.

Carbolic Acid (No. 2 Calvert) 20 grs.

Hot Water 1 fl. oz.

Dissolve.

INHALATIO ACIDI SULPHUROSII.

Inhalation of Sulphurous Acid.

Sulphurous Acid	3 fl. drs.
Water	to 1 fl. oz.

Mix.

INHALATIO ACIDI TANNICI.

Inhalation of Tannic Acid.

Tannic Acid	10 grs.
Water	1 fl. oz.

Dissolve.

INHALATIO ALKALINA.

Alkaline Inhalation.

Carbonate of Soda	10 grs.
Water	1 fl. oz.

Dissolve.

INHALATIO ALUMINIS.

Inhalation of Alum.

Alum	15 grs.
Water	1 fl. oz.

Dissolve.

INHALATIO POTASSÆ CHLORATIS.

Inhalation of Chlorate of Potash.

Chlorate of Potash	to saturation.
Water	1 fl. oz.

Dissolve.

Juice, Todi 3i

Water of in Mau's Inhaler

Spray for Diphtheria

Carbolic Acid gr 120

Todin Lin.

$\frac{3 \text{ } \overline{\text{iv}}}{\text{}}$

Recs. Spirit

$\frac{3 \text{ } \overline{\text{i}}}{\text{}}$

Aq ad

$\frac{3 \text{ } \times \text{II}}{\text{}}$

4th.

INJECTIO CURARE HYPODERMICA.

Hypodermic Injection of Curare.

Curare 4 grs.
 Distilled Water 1 fl. dr.

Rub the Curare with the Water, allow it to stand for 48 hours, and then filter.

INJECTIO ERGOTINÆ HYPODERMICA.

Hypodermic Injection of Ergotine.

Ergotine (Bonjean's) 10 grs.
 Distilled Water 1 fl. dr.

Dissolve.

INJECTIO MORPHIÆ HYPODERMICA.

Hypodermic Injection of Morphia.

Made of the same strength as the Injectio Morphiæ Hypodermica, B. P., by dissolving freshly prepared Acetate of Morphia in distilled water.

TABLE OF STRENGTH.

1 grain of Acetate of Morphia in 12 minims.

$\frac{3}{4}$	"	"	"	9	"
$\frac{2}{3}$	"	"	"	8	"
$\frac{1}{2}$	"	"	"	6	"
$\frac{1}{3}$	"	"	"	4	"
$\frac{1}{4}$	"	"	"	3	"
$\frac{1}{6}$	"	"	"	2	"
$\frac{1}{9}$	"	"	"	$1\frac{1}{2}$	"

IODIFORMUM.

Iodoform.

Used in the form of powder, in the Ointment and in the Ethereal Solution of Iodoform.

LAPIS DIVINUS.

"Pierre Divine."

Sulphate of Copper	} of each	. 3 oz.
Nitrate of Potash		
Alum		
Camphor		60 grs.

Rub the Copper, Potash, and Alum into a powder and heat them in a crucible until fused; then add the Camphor in fine powder. Cast into moulds, and keep in a stoppered bottle.

LINCTUS OPIATUS.

Opiate Linctus.

Tincture of Opium	10 m.	<i>(1 1/4 to 1 1/2)</i>
Dilute Sulphuric Acid	10 m.	<i>dose of 3i.</i>
Treacle	6 drs.	
Water	to 1 fl. oz.	

Mix the Treacle with the Water, then add the Acid and Tincture.

Dose.—1 to 2 *teaspoonfuls*.

Boracic & Iodoform Emulsion

Acidi Borici gr. $\overline{\text{X}}$

Iodoform $\overline{\text{ʒ}}\text{ss}$

Glycerini $\overline{\text{ʒ}}\text{ij}$

Pulv. Tragacanth Co $\overline{\text{ʒ}}\text{ss}$

Aqua $\overline{\text{ʒ}}\text{viij}$

1 Part in 8 to be used

Hair. Wash

Sodae Bicarb	3 ⁷ / ₈
Sp. Rect	3 ¹¹ / ₁₆
Liq Am. B. P.	3 ¹⁴ / ₁₆
Sp. Rosmar	3 ¹ / ₁₆
Aq Camph ad	3 ^X / ₁₆

LINIMENTUM MYRISTICÆ.

Liniment of Nutmeg.

Expressed Oil of Nutmeg	.	.	.	.	1 oz.
Olive Oil	.	.	.	.	3 fl. oz.

Mix.

LINIMENTUM SAPONIS MOLLIS.

Liniment of Soft Soap.

Soft Soap	.	.	.	.	12 oz.
Camphor	.	.	.	.	4 oz.
Oil of Origanum	.	.	.	.	1 fl. oz.
Rectified Spirit	.	.	.	.	3 pints.

Put the Soap into the Spirit, and place them in a warm place. Stir frequently. When dissolved, add the Oil and Camphor.

LINTEUM BORACICUM.

Boracic Lint.

Lint steeped in a hot saturated solution of Boracic Acid, and dried without wringing.

“LIQUOR CARBONIS DETERGENS.”

Used in the Lotion.

LIQUOR IODOFORMI ÆTHEREUS.

Ethereal Solution of Iodoform.

A saturated solution of Iodoform in Ether.
For local application.

LOTIO ACIDI CARBOLICI.

Carbolic Acid Lotion. (1 in 20.)

Carbolic Acid	.	.	.	.	24	grs.
Water	.	.	.	.	1	fl. oz.

Dissolve.

LOTIO ACIDI NITRICI.

Nitric Acid Lotion.

Dilute Nitric Acid	.	.	.	.	10	m.
Water	.	.	.	.	to 1	fl. oz.

Mix.

LOTIO ACIDI HYDROCYANICI.

Hydrocyanic Acid Lotion.

Dilute Hydrocyanic Acid	.	.	.	.	10	m.
Distilled Water	.	.	.	.	to 1	fl. oz.

Mix.


LOTIO ACIDI TANNICI.

Tannic Acid Lotion.

Tannic Acid	.	.	.	.	10	grs.
Distilled Water	.	.	.	.	1	fl. oz.

Dissolve.

To dissolve wash in the ear
Liq. Potassae 3ij to ops warm water



LOTIO ALKALINA.

Alkaline Lotion.

Carbonate of Soda	.	.	.	7 grs.
Water	.	.	.	1 fl. oz.

Dissolve.

LOTIO ALUMINIS.

Alum Lotion.

Alum	.	.	.	.	10 grs.
Water	.	.	.	.	1 fl. oz.

Dissolve.

LOTIO ALUMINIS ET ZINCI
SULPHATIS.*Alum and Zinc Lotion.*

Alum	.	.	.	.	10 grs.
Sulphate of Zinc	.	.	.	.	10 grs.
Distilled Water	.	.	.	.	1 fl. oz.

Dissolve.

LOTIO ARGENTI NITRATIS.

Nitrate of Silver Lotion.

Nitrate of Silver	.	.	.	20 grs.
Distilled Water	.	.	.	4 fl. oz.

Dissolve.

LOTIO BORACIS.

33 $\frac{1}{3}$ in 2*Borax Lotion.*

Borax	.	.	.	.	.	10 grs.
Distilled Water	.	.	.	.	.	1 fl. oz.

Dissolve.

LOTIO LIQUORIS CARBONIS DETERGENTIS.

Spirit of Coal Tar Lotion.

"Liquor Carbonis Detergens"	.	.	.	.	.	$\frac{1}{2}$ to 1 fl. oz.
Water	.	.	.	.	.	to 1 pint.

Mix.

LOTIO GLYCERINI ALKALINA.

Alkaline Glycerine Lotion.

Glycerine	.	.	.	.	.	1 fl. dr.
Carbonate of Soda	.	.	.	.	.	5 grs.
Distilled Water	.	.	.	.	.	to 1 fl. oz.

Mix.

LOTIO HYDRARGYRI PERCHLORIDI.

Perchloride of Mercury Lotion.

Perchloride of Mercury	.	.	.	.	.	2 grs.
Distilled Water	.	.	.	.	.	1 fl. oz.

Dissolve

Thompson's Fluid

Borax $\overline{3\text{ i}}$

Glycerine $\overline{3\text{ ii}}$

Aqua $\overline{3\text{ ii}}$ $\overline{3\text{ iv}}$ to the vj .

Mouth Wash.

Glyc. Boracis $\overline{3\text{ ii}}$


Succus Limonis $\overline{3\text{ ii}}$

Aqua ad $\overline{3\text{ i}}$

Salicylic Cream.

S. Acid 1 part

Vaseline 8 parts



LOTIO IODI.

Iodine Lotion.

Tincture of Iodine	.	.	.	20 m.
Distilled Water	.	.	.	to 1 fl. oz.

Mix.

LOTIO OPII.

Opium Lotion.

Extract of Opium	.	.	.	5 grs.
Water	.	.	.	1 fl. oz.

Mix.

LOTIO PLUMBI VEL ALBA.

"Goulard Wash."

The "Diluted Solution of Subacetate of Lead."
B. P.

LOTIO PLUMBI ACETATIS CUM
MORPHIA.*Lotion of Acetate of Lead with Morphia.*

Acetate of Lead	.	.	.	1 gr.
Acetate of Morphia	.	.	.	1 gr.
Distilled Water	.	.	.	1 fl. oz.

Dissolve.

LOTIO PLUMBI CUM OPIO.

"Goulard Wash," with Opium.

"Goulard Wash"	}	. . .	equal parts.
Lotion of Opium			

Mix.

LOTIO POTASSÆ CHLORATIS.

Chlorate of Potash Lotion.

Chlorate of Potash	.	.	.	10 grs.
Water	.	.	.	1 fl. oz.

Dissolve.

LOTIO POTASSÆ PERMANGANATIS.

Permanganate of Potash Lotion.

Solution or Permanganate of				
Potash	.	.	.	10 m.
Distilled Water	.	.	.	1 fl. oz.

Mix.

LOTIO SODÆ CHLORINATÆ.

Chlorinated Soda Lotion.

Solution of Chlorinated Soda	.	1 fl. dr.
Distilled Water	.	1 fl. oz.

Mix.

This may be used as a gargle.

LOTIO SODÆ HYPOSULPHITIS.

Hyposulphite of Soda Lotion.

Hyposulphite of Soda	.	.	.	1 dr.
Distilled Water	.	.	.	1 fl. oz.

Dissolve.

LOTIO SPIRITUS.

Spirit Lotion.

Rectified Spirit	.	.	.	2 fl. drs.
Distilled Water	.	.	.	to 1 fl. oz.

Mix.

LOTIO THYMOL.

Thymol Lotion. (1 in 1000.)

Thymol	.	.	.	.	10 grs.
Hot Water	.	.	.	.	1 pint.

Dissolve.

LOTIO ZINCI OXIDI.

Oxide of Zinc Lotion.

Oxide of Zinc	.	.	.	.	$\frac{1}{2}$ dr.
Glycerine	.	.	.	.	1 fl. dr.
Distilled Water	.	.	.	.	to 1 fl. oz.

Mix.

LOTIO ZINCI SULPHATIS.

Sulphate of Zinc Lotion.

Sulphate of Zinc 2 grs.

Distilled Water 1 fl. oz.

Dissolve.

LOTIO ZINCI SULPHATIS ET ACIDI CARBOLICI.

Lotion of Sulphate of Zinc and Carbolic Acid.

Sulphate of Zinc 2 grs.

Carbolic Acid 10 grs.

Distilled Water to 1 fl. oz.

Dissolve.

MISTURA ACIDI HYDROCHLORICI.

Mixture of Hydrochloric Acid.

Dilute Hydrochloric Acid . . . 10 m.

Simple Syrup 1 fl. dr.

Distilled Water to 1 fl. oz.

Mix.

Dose.—1 to 2 fl. oz.

MISTURA ACIDI NITRICI.

Nitric Acid Mixture.

Dilute Nitric Acid 10 m.

Simple Syrup 1 fl. dr.

Distilled Water to 1 fl. oz.

Mix.

Dose.—1 to 2 fl. oz.

Q.
Liq. Ext. Case. Sag: $\eta \overline{X}$ (to $\eta \overline{XXX}$)
Mist. Acidi Co. ad $\overline{31}$. f. o.

MISTURA ACIDI NITRO-HYDRO- CHLORICI.

Mixture of Nitro-Hydrochloric Acid.

Dilute Nitro-Hydrochloric Acid .	10 m.
Simple Syrup	1 fl. dr.
Distilled Water	to 1 fl. oz.

Mix.

Dose.—1 to 2 fl. oz.

MISTURA ACIDI COMPOSITA.

*Compound Mixture of Nitro-Hydrochloric
Acid.*

Tincture of Nux Vomica . . .	10 m.
Compound Tincture of Gentian .	1 fl. dr.
Mixture of Nitro-Hydrochloric Acid	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA ALUMINIS COMPOSITA.

Compound Mixture of Alum.

Alum	8 grs.
Compound Tincture of Camphor .	20 m.
Ipecacuanha Wine	15 m.
Camphor Water	$\frac{1}{2}$ fl. oz.
Dill Water	to 1 fl. oz.

Mix.

Dose.—For a child, 1 to 2 fl. drs.; for an adult,
1 fl. oz.

MISTURA AMMONIÆ.

"Ammonia Julep."

Carbonate of Ammonia	. . .	4 grs.
Treacle		20 m.
Compound Tincture of Lavender		20 m.
Peppermint Water		to 1 fl. oz.

Dissolve the Treacle in the Peppermint Water,
then add the Ammonia and Lavender.

Dose.—1 to 1½ fl. oz.

MISTURA AMMONIÆ ACETATIS.

"Diluted Solution of Acetate of Ammonia."

Solution of Acetate of Ammonia	} equal parts.
Distilled Water	

Mix.

Dose.—1 fl. oz.

MISTURA AMMONIÆ EFFERVESCENS.

Effervescing Mixture of Ammonia.

Carbonate of Ammonia		20 grs.
Distilled Water		1 fl. oz.

Dissolve.

Lemon Juice ½ fl. oz.
The solution of Ammonia and Lemon Juice to be
put in separate bottles.

Dose.—1 fl. oz. of the former and ½ fl. oz. of the
latter to be mixed and given in an effervescing
state.

MISTURA ANODYNA.

Anodyne Mixture.

Spirit of Camphor	.	.	.	$\frac{1}{2}$ fl. dr.
Tincture of Hyoscyamus	.	.	.	$\frac{1}{2}$ fl. dr.
Mucilage of Tragacanth	.	.	.	$\frac{1}{2}$ fl. dr.
Distilled Water	.	.	.	to 1 fl. oz.

Mix.

Dose.—1 to $1\frac{1}{2}$ fl. oz.

MISTURA BALSAMI PERUVIANI.

Mixture of Balsam of Peru.

Balsam of Peru	.	.	.	25 m.
Unclarified Honey	.	.	.	90 grs.
Distilled Water	.	.	.	to 1 fl. oz.

Rub the Balsam of Peru with the Honey ; when well mixed, gradually add the Water.

Mix.

Dose.—1 fl. oz.

MISTURA BELLADONNÆ COMPOSITA.

Compound Mixture of Belladonna.

Tincture of Belladonna	.	.	.	10 m.
Dilute Hydrocyanic Acid	.	.	.	2 m.
Bicarbonate of Soda	.	.	.	10 grs.
Simple Syrup	.	.	.	1 fl. dr.
Distilled Water	.	.	.	to 1 fl. oz.

Mix.

Dose.—For a child, 1 to 2 fl. drs. ; for an adult, 1 fl. oz.

MISTURA BISMUTHI.

Bismuth Mixture.

Subnitrate of Bismuth. . . .	10 grs.
Compound Powder of Tragacanth	10 grs.
Distilled Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA BISMUTHI SEDATIVA.

Sedative Mixture of Bismuth.

Subnitrate of Bismuth. . . .	10 grs.
Bicarbonate of Soda	10 grs.
Solution of Hydrochlorate of Mor- phia	10 m.
Compound Powder of Tragacanth	10 grs.
Distilled Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA CASCARILLÆ COMPOSITA.

Compound Mixture of Cascarilla.

Tincture of Squills	15 m.
Compound Tincture of Camphor	1 fl. dr.
Infusion of Cascarilla	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

Ac. Hydrocyanic ℥i ℥iv
Mist. Bismuthi ℥i .
(For indigestion)

MISTURA CINCHONINÆ.

Cinchonine Mixture.

Hydrochlorate of Cinchonine	.	2	grs.
Dilute Hydrochloric Acid	.	4	m.
Tincture of Orange	.	10	m.
Simple Syrup	.	$\frac{1}{2}$	fl. dr.
Distilled Water	.	to 1	fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA CONII COMPOSITA.

Compound Hemlock Mixture.

Extract of Hemlock	.	5	grs.
Bicarbonate of Soda	.	5	grs.
Spirit of Nutmeg	.	$\frac{1}{2}$	fl. dr.
Liquorice Water	.	to 1	fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA COPAIBÆ.

Copaiva Mixture.

Copaiva	.	20	m.
Tincture of Quillaia	.	20	m.
Tincture of Lemon Peel	.	20	m.
Water	.	to 1	fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA COPAIBÆ RESINÆ.

Mixture of Resin of Copaiva.

Resin of Copaiva	15 grs.
Rectified Spirit of Wine . .	20 m.
Compound Powder of Tragacanth	15 grs.
Syrup of Ginger	1 fl. dr.
Distilled Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA CRETÆ COMPOSITA.

“Diarrhœa Mixture.”

Aromatic Spirit of Ammonia .	20 m.
Tincture of Catechu	20 m.
Mixture of Chalk	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.


MISTURA ERGOTÆ.

Ergot Mixture.

Liquid Extract of Ergot . .	$\frac{1}{2}$ fl. dr.
Tincture of Orange Peel . .	$\frac{1}{2}$ fl. dr.
Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.



R. Tr Ferri Perchlor
 Tr. Nucis Vom.
 Acid Phos. Dil
 Spt. Armoracia
 Glycerini
 Aqua ad

℥i
 ℥i
 ℥viii
 3℥
 3℥
 3i.

F. & S.

for throat.

R. Ferri Perchlor
 Mist. Acidi Co

℥i

3i. F. & S. p.c

Spray for tonsillitis.

Boracis 3i

(M. Durham)

Acidi Borici 3℥

Cocain. Hydr. gr XII

Aqua ^{distil} ad 3vi. 3℥ss. in Syring.

Gargle, after extraction of tonsils

Borax 3i

Acidi Borici 3℥

Liq. Morph. Hydr 3iv

Aqua ad 3viii.

MISTURA FERRI TARTARATI.

Mixture of Tartarated Iron.

Tartarated Iron	10 grs.
Rectified Spirit of Wine	$\frac{1}{2}$ fl. dr.
Simple Syrup	1 fl. dr.
Water	to 1 fl. oz.

Mix.

Dose.—*For a child, 1 to 4 fl. drs.; for an adult,*
1 fl. oz.

MISTURA FERRI ET QUASSIÆ.

Mixture of Iron and Quassia.

Tincture of Perchloride of Iron	15 m.
Infusion of Quassia	to 1 fl. oz.

Mix.

Dose.—*1 fl. oz.*

MISTURA FILICIS MARIS.

Mixture of Male Fern.

Liquid Extract of Male Fern	1 fl. dr.
Syrup of Ginger	$\frac{1}{2}$ fl. dr.
Tincture of Quillaia	$\frac{1}{2}$ fl. dr.
Water	to 1 fl. oz.

Mix.

Dose.—*1 fl. oz.*

MISTURA GENTIANÆ COMPOSITA.

Compound Mixture of Gentian.

Compound Tincture of Carda-	
moms	80 m.
Compound Infusion of Gentian	$\frac{1}{2}$ fl. oz.
Infusion of Senna	$\frac{1}{2}$ fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA HÆMATOXYLI COMPOSITA.

Compound Mixture of Logwood.

Extract of Logwood, in powder	10 grs.
Ipecacuanha Wine	10 m.
Opium Wine	5 m.
Chalk Mixture	to 1 fl. oz.

Rub the Extract of Logwood with the Chalk
Mixture until well mixed, then add the Wines.

Dose.—1 fl. oz.

MISTURA HYDRARGYRI
PERCHLORIDI.*Mixture of Perchloride of Mercury.*

Solution of Perchloride of Mercury	80 m.
Infusion of Quassia	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

Vini Colchici. vj I

Por. Baccar gr I ss

Mist Gent Co ad 3i .

(For Lumbago).

Hemorrhage fr Lungs

Opopentum vj ss

Brown sugar

Yolk of egg. _____

MISTURA IODI.

Iodine Mixture.

Iodine	1/8 gr.
Iodide of Potassium	3 grs.
Peppermint Water	to 1 fl. oz.

Dissolve.

Dose.—1 fl. oz.

MISTURA KRAMERIÆ.

Rhatany Mixture.

Ipecacuanha Wine	20 m.
Tincture of Catechu	20 m.
Infusion of Rhatany Root	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA MAGNESIÆ.

Mixture of Carbonate of Magnesia.

Carbonate of Magnesia	15 grs.
Water	1 fl. oz.

Mix well together in a mortar.

Dose.—1 fl. oz.

MISTURA MAGNESIÆ ET MAGNESIÆ SULPHATIS.

"White Mixture."

Carbonate of Magnesia	. . .	10 grs.
Sulphate of Magnesia	. . .	1 dr.
Peppermint Water	. . .	1 fl. oz.

Dissolve the Sulphate of Magnesia in the Peppermint Water, and finally rub down with the Carbonate in a mortar.

Dose.—1 *fl. oz.*

MISTURA MUCILAGINOSA.

Mucilage Mixture.

Mucilage of Gum Acacia	. . .	2½ fl. dr.
Almond Oil	. . .	10 m.
Syrup	. . .	½ fl. dr.
Water	. . .	to 1 fl. oz.

Mix well the Oil with the Mucilage, then add the Syrup; these being well mixed, add the Water gradually.

Dose.—1 *fl. oz.*

MISTURA OLEI.

Olive Oil Mixture.

Olive Oil	. . .	1 fl. dr.
Solution of Potash	. . .	4 m.
Distilled Water	. . .	to 1 fl. oz.

Mix.

Dose.—1 *fl. oz.*

MISTURA OLEI MORRHUÆ ET FERRI.

Mixture of Cod Liver Oil and Iron.

Cod Liver Oil	$\frac{1}{2}$ fl. oz.
Tartarated Iron	10 grs.
Solution of Potash	3 m.
Water	to 1 fl. oz.

Shake well together until they blend.

Dose.—2 to 4 *fl. drs.* .

MISTURA OLEI RICINI.

Castor Oil Mixture.

Castor Oil	1 fl. dr.
Tincture of Rhubarb	1 fl. dr.
Tincture of Quillaia	20 m.
Pimento Water	to 1 fl. oz.

Mix.

Dose.—1 *fl. oz.*

MISTURA OLIBANI.

Olibanum Mixture.

Olibanum, powdered	20 grs.
Honey	$\frac{1}{2}$ fl. dr.
Barley Water	to 1 fl. oz.

Rub thoroughly the Olibanum with the Honey in a mortar, then gradually add the Barley Water.

Dose.—1 *fl. oz.*

MISTURA OPII ACIDA.

Acid Opium Mixture.

Tincture of Opium	.	.	.	5 m.
Aromatic Sulphuric Acid	.	.	.	10 m.
Syrup of Tolu	.	.	.	1 fl. dr.
Water	.	.	.	to 1 fl. oz.

Mix.

Dose.—*To 1 fl. oz.*

MISTURA OXYMELLIS.

Oxymel Mixture.

Oxymel	.	.	.	.	3 fl. drs.
Nitrate of Potash	.	.	.	.	7 grs.
Water	.	.	.	.	to 1 fl. oz.

Mix.

Dose.—*1 fl. oz.*

MISTURA OXYMELLIS COMPOSITA.

Compound Mixture of Oxymel.

Ipecacuanha Wine	.	.	.	5 m.
Compound Tincture of Camphor.	.	.	.	20 m.
Oxymel Mixture	.	.	.	to 1 fl. oz.

Mix.

Dose.—*1 fl. oz.*

Tr. Camph. Co
 Syrupi Tolu
 Myrtillicis Scilla } (Mistura Swayne)
 a a ad 3i. l. ss.

Liq. Morph. Bimec. 3ij
 Tinct. Sassa 3p.
 Syr. Codeia 3ij.
 Aq ad 3ij
 3i occasionally when
 the cough is troublesome.

Vin. Specac 7ij
 Syr. Tolu 3i
 Tr. Camph. Co 3p
 Acid. Hydrochlor. Dil 7ij
 Aqua Chloroformi ad 3i.
 w. be for first 2 hrs.
 after that w. 2 hrs.

MISTURA PAREIRÆ COMPOSITA.

Compound Mixture of Pareira.

Bicarbonate of Potash . . .	20 grs.
Tincture of Hyoscyamus . . .	$\frac{1}{2}$ fl. dr.
Decoction of Pareira . . .	to 1 fl. oz.

Mix.

Dose.—1 to $1\frac{1}{2}$ fl. oz.

MISTURA POTASSÆ ACETATIS COMPOSITA.

“Diuretic Mixture.”

Acetate of Potash . . .	20 grs.
Tincture of Squills . . .	15 m.
Spirit of Nitrous Ether . . .	$\frac{1}{2}$ fl. dr.
Scoparium Juice . . .	1 fl. dr.
Distilled Water . . .	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA POTASSÆ CHLORATIS.

Chlorate of Potash Mixture.

Chlorate of Potash, in powder . .	10 grs.
Decoction of Cinchona . . .	1 fl. oz.

Dissolve.

Dose.—1 fl. oz.

MISTURA POTASSÆ CITRATIS.

"Saline Mixture."

Citrate of Potash	20 grs.
Water	1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA POTASSÆ EFFERVESCENS.

Effervescing Mixture of Potash.

Bicarbonate of Potash	20 grs.
Distilled Water	to 1 fl. oz.

Mix.

Lemon Juice	$\frac{1}{2}$ fl. oz.
-----------------------	-----------------------

To be taken in an effervescing state.

Dose.—1 fl. oz. of the solution of Bicarbonate of Potash to $\frac{1}{2}$ fl. oz. of the Lemon Juice.

MISTURA POTASSÆ NITRATIS.

Nitrate of Potash Mixture.

Nitrate of Potash	15 grs.
Spirit of Nitrous Ether	15 m.
Syrup of Lemons	40 m.
Peppermint Water	to 1 fl.

Mix.

Dose.—1 fl. oz.

MISTURA POTASSII IODIDI.

Iodide of Potassium Mixture.

Iodide of Potassium	5 grs.
Aromatic Spirit of Ammonia	20 m.
Compound Infusion of Gentian	4 fl. drs.
Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA QUININÆ.

Quinine Mixture.

Sulphate of Quinine	2 grs.
Dilute Sulphuric Acid.	4 m.
Tincture of Orange Peel	10 m.
Simple Syrup	$\frac{1}{2}$ fl. dr.
Water	to 1 fl. oz.

Dissolve the Quinine in the Acid, first mixed in $\frac{1}{2}$ oz. of the Water, then add the other ingredients, and the Water.

Dose.—1 to $1\frac{1}{2}$ fl. oz.

MISTURA RHEI COMPOSITA.

Compound Rhubarb Mixture.

Rhubarb Root, in powder	5 grs.
Calumba Root, in powder	10 grs.
Bicarbonate of Soda	10 grs.
Peppermint Water	1 fl. oz.

Rub the Rhubarb, Calumba, and Soda well together, then gradually add the Peppermint Water.

Dose.—1 fl. oz.

MISTURA ROSÆ LAXATIVA.

Laxative Mixture of Roses.

Sulphate of Magnesia . . .	40 grs.
Pimento Water . . .	2 fl. drs.
Acid Infusion of Roses . . .	to 1 fl. oz.

Dissolve.

Dose.—1 fl. oz.

MISTURA SENEGÆ.

Senega Mixture.

Carbonate of Ammonia . . .	5 grs.
Tincture of Senega . . .	$\frac{1}{2}$ fl. dr.
Infusion of Senega . . .	$\frac{1}{2}$ fl. oz.
Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

MISTURA SENNÆ.

“House Medicine.”

Sulphate of Magnesia . . .	1 dr.
Infusion of Senna . . .	1 fl. oz.

Mix.

Dose.—1 to 2 fl. oz.

MISTURA SEVI.

Suet and Milk Mixture.

Mutton Suet, cut very fine . . .	1 oz.
Water	4 fl. oz.

Boil gently for ten minutes, and strain; then add

New Milk	16 fl. oz.
Cinnamon, slightly bruised . . .	60 grs.
Refined Sugar	1 oz.

Boil again for ten minutes, and strain.

Dose.—4 to 8 oz.

MISTURA TEREBINTHINÆ.

Turpentine Mixture.

Oil of Turpentine	20 m.
Tincture of Quillaia	20 m.
Tincture of Lemon Peel	$\frac{1}{2}$ fl. dr.
Water	to 1 fl. oz.

Mix.

Dose.—1 fl. oz.

OLEUM ACIDI CARBOLICI.

Carbolic Acid Oil. (1 in 20.)

Carbolic Acid	24 grs.
Olive Oil	1 fl. oz.

Mix.

PASTA CAUSTICA.

Caustic Paste.

Extract of Sanguinaria	.	.	$\frac{1}{2}$ oz.
Extract of Stramonium	.	.	1 oz.
Mix, and add when wanted			
Chloride of Zinc	.	.	$\frac{1}{2}$ oz.

PASTA CAUSTICA CUM OPIO.

Caustic Paste with Opium.

Fused Chloride of Zinc	.	.	480 grs.
Flour	.	.	180 grs.
Battley's Solution of Opium	.	.	1 fl. oz.
Mix until a paste is formed.			

PILULÆ ACIDI CARBOLICI
COMPOSITÆ.*Compound Carbolic Acid Pills.*

Carbolic Acid (No. 1, Calvert)	.	.	12 grs.
Guaiacum Resin	.	.	12 grs.
Powdered Rhubarb	.	.	24 grs.

Make 12 pills.

Dose.—1 pill.


PILULÆ ANTIMONII OPIATÆ.

Antimony and Opium Pills.

Tartarated Antimony	.	.	3 grs.
Opium, in powder	.	.	6 grs.
Treacle	.	.	a sufficiency.

Make 12 pills.

Dose.—1 pill.



Laxative Pill for chronic.

℞ Nucis Vom gr $\dot{\text{r}}$..
℞ Aloes (Barb) gr $\dot{\text{r}}$..
℞ Hyoscy. gr $\dot{\text{r}}$..

or
℞ Belladonna } a. a. gr $\frac{1}{4}$
℞ Nucis Vom } gr $\dot{\text{r}}$
℞ Aloes } gr $\dot{\text{r}}$
Pile Rhei Co. gr $\dot{\text{r}}$

1. Lds.

PILULÆ CALOMELANOS CUM OPIO.

Calomel and Opium Pills.

Calomel 12 grs.

Opium 3 grs.

Confection of Roses . . . a sufficiency.

Make 12 pills.

Dose.—1 pill.

PILULÆ CAMPHORÆ ET HYOSCYAMI.

Camphor and Henbane Pills.

Camphor 24 grs.

Rectified Spirit 4 m.

Rub into a fine powder, then add

Extract of Hyoscyamus . . . 36 grs.

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ COLOCYNTHIDIS CUM
CALOMELANE.*Colocynth and Calomel Pills.*

Compound Pill of Colocynth . 48 grs.

Subchloride of Mercury . . 12 grs.

Make 12 pills.

Dose.—1, 2, or 3 pills.

PILULÆ CONII ET ACIDI BENZOICI.

Hemlock and Benzoic Acid Pills.

Extract of Conium	.	.	.	24	grs.
-------------------	---	---	---	----	------

Benzoic Acid	.	.	.	36	grs.
--------------	---	---	---	----	------

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ CONII ET BISMUTHI.

Hemlock and Bismuth Pills.

Extract of Hemlock	.	.	.	30	grs.
--------------------	---	---	---	----	------

Subnitrate of Bismuth	.	.	.	40	grs.
-----------------------	---	---	---	----	------

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ CONII ET HYDRARGYRI.

Hemlock and Mercury Pills.

Extract of Conium	.	.	.	50	grs.
-------------------	---	---	---	----	------

Mercurial Pill	.	.	.	12	grs.
----------------	---	---	---	----	------

Make 12 pills.

Dose.—1 pill.

PILULÆ CREASOTI.


Creasote Pills.

Creasote	.	.	.	.	12	m.
----------	---	---	---	---	----	----

Yellow Wax	.	.	.	.	36	grs.
------------	---	---	---	---	----	------

Make 12 pills.

Dose.—1 pill.



PILULÆ CUPRI SULPHATIS.

Sulphate of Copper Pills.

Sulphate of Copper	.	.	.	3 grs.
Opium, in powder	.	.	.	6 grs.
Confection of Hips	.	.	.	a sufficiency.

Make 12 pills.

Dose.—1 *pill*.PILULÆ DIURETICÆ CUM
HYDRARGYRO.*Diuretic Pills with Mercury.*

Mercurial Pill	.	.	.	12 grs.
Squill Root, in powder	.	.	.	12 grs.
Digitalis Leaf, in powder	.	.	.	12 grs.
Extract of Hyoscyamus	.	.	.	20 grs.

Make 12 pills.

Dose.—1 *pill*.

PILULÆ GENTIANÆ ET FERRI.

Gentian and Iron Pills.

Extract of Gentian	.	.	.	36 grs.
Sulphate of Iron	.	.	.	12 grs.

Make 12 pills.

Dose.—1 or 2 *pills*.

PILULÆ GENTIANÆ ET ZINCI.

Gentian and Zinc Pills.

Extract of Gentian	.	.	.	36 grs.
Sulphate of Zinc	.	.	.	12 grs.
Calumba Root, in powder	.			a sufficiency.

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ HYDRARGYRI ET OPII.

Mercury and Opium Pills.

Mercurial Pill	.	.	.	48 grs.
Opium, in powder	.	.	.	3 grs.

Make 12 pills.

Dose.—1 pill.

PILULÆ HYDRARGYRI ET RHEI.

Mercury and Rhubarb Pills.

Mercurial Pill	.	.	.	30 grs.
Compound Rhubarb Pill	.	.	.	30 grs.

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ HYOSCYAMI ET MORPHIÆ.

Henbane and Morphia Pills.

Extract of Hyoscyamus	.	.	36 grs.
Hydrochlorate of Morphia	.	.	4 grs.

Make 12 pills.

Dose.—1 pill.

PILULÆ IPECACUANHÆ COMPOSITÆ.

“Dover's Pills.”

Compound Ipecacuanha Powder . 60 grs.
Treacle a sufficiency.

Make 12 pills.

Dose.—1 or 2 pills.

PILULÆ NUCIS VOMICÆ.

Nux Vomica Pills.

Extract of Nux Vomica . . 6 grs.
Powdered Capsicum . . 3 grs.
Compound Rhubarb Pill . . 24 grs.

Make 12 pills.

Dose.—1 pill.

PILULÆ NUCIS VOMICÆ ET FERRI.

Nux Vomica and Iron Pills.

Extract of Nux Vomica . . 6 grs.
Dried Sulphate of Iron . . 12 grs.
Extract of Barbadoes Aloes . . 12 grs.

Make 12 pills.

$2\frac{1}{2}$ grs. Dose.—1 pill.

PILULÆ PODOPHYLLI CUM
HYDRARGYRO.

Podophyllum and Mercury Pills.

Resin of Podophyllum	.	.	6 grs.
Mercurial Pill	.	.	12 grs.
Extract of Hyoscyamus	.	.	24 grs.

Make 12 pills.

Dose.—1 *pill*.

PILULÆ SCILLÆ ET HYDRARGYRI.

Squill and Mercury Pills.

Grey Oxide of Mercury	.	.	4 grs.
Compound Squill Pill	.	.	48 grs.

Make 12 pills.

Dose.—1, 2, or 3 *pills*.

PILULÆ SCILLÆ ET OPII.

Squill and Opium Pills.

Opium, in powder	.	.	4 grs.
Compound Squill Pill	.	.	48 grs.

Make 12 pills.

Dose.—1, 2, or 3 *pills*.

PILULÆ THYMOL.

Thymol Pills.

Thymol	.	.	.	.	.	12	grs.
Gum Tragacanth	.	.	.	.	.	12	grs.
Mastich	.	.	.	.	.	12	grs.
Glycerine	.	.	.	.	.	a sufficiency.	

Make 12 pills.

Dose.—1 or 2 pills.

PULVIS ELATERII SALINUS.

Saline Powder of Elaterium.

Extract of Elaterium, in powder	.	.	.	.	1	gr.
Acid Tartrate of Potash	.	.	.	.	25	grs.
Ginger, in powder	.	.	.	.	4	grs.

Mix thoroughly.

Dose.—5 to 30 grs.

PULVIS EMETICUS.

Emetic Powder.

Ipecacuanha Root, in powder	.	.	.	.	20	grs.
Tartarated Antimony	.	.	.	.	$\frac{1}{2}$	gr.

Mix.

Dose.—5 to 20 grs.

PULVIS JALAPÆ CUM CALOMELANE.

Powder of Jalap and Calomel.

Jalap, in powder . . .	20 grs.
Subchloride of Mercury . . .	5 grs.
Ginger, in powder . . .	5 grs.

Mix.

Dose.—*From 10 to 30 grs.*

PULVIS RHEI ET SODÆ.

Powder of Rhubarb and Soda.

Rhubarb, in powder . . .	5 grs.
Carbonate of Soda, dried . . .	5 grs.
Calumba, in powder . . .	10 grs.

Rub well together.

Dose.—*10 to 20 grs.*

PULVIS RHEI SALINUS.

Saline Rhubarb Powder.

Rhubarb, in powder . . .	10 grs.
Sulphate of Potash, in powder . . .	20 grs.

Mix.

Dose.—*From 15 to 40 grs.*

PULVIS SALIUM.

Powder of Salts.

Sulphate of Soda	30 grs.
Sulphate of Magnesia	30 grs.
Chloride of Soda	1 gr.
Bicarbonate of Soda	1 gr.

Dose.—1 to 4 *drs.*PULVIS SCAMMONII CUM
CALOMELANE.*Powder of Scammony and Calomel.*

Scammony, in powder	6 grs.
Subchloride of Mercury	3 grs.
Refined Sugar	3 grs.

Mix.

Dose.—10 to 20 *grs.*

PULVIS SODÆ ET HYDRARGYRI.

Powder of Soda and Mercury.

Carbonate of Soda, dried	5 grs.
Subchloride of Mercury	1 gr.
Aromatic Powder of Chalk	10 grs.

Rub well together.

Dose.—From 4 to 16 *grs.*

PULVIS SODÆ ET HYDRARGYRI CUM CRETA.

Powder of Soda and Mercury with Chalk.

Carbonate of Soda . . . 3 grs.

Mercury with Chalk . . . 1 gr.

Mix.

Dose.—*From 4 to 10 grs.*

QUILLAIÆ CORTEX.

Bark of the Quillaia Saponaria.

Used in the Tincture of Quillaia Bark.

THYMOL.

A Crystalline Substance of the Phenol Series, occurring naturally in the Oil of Thyme.

Used in the Lotion and Pill of Thymol.

TINCTURA CINCHONÆ COMPOSITA.

Compound Tincture of Cinchona.

The Compound Tincture of Cinchona (B.P.), omitting the Saffron.

TINCTURA PODOPHYLLI RESINÆ.

Tincture of Podophyllum Resin.

Resin of Podophyllum . . . 8 grs.

Rectified Spirit of Wine . . . 1 fl. oz.

Dissolve.

Dose.—*20 m. to 1 fl. dr.*

TINCTURA QUILLAIÆ.

Tincture of Quillaia Bark.

Quillaia Bark, in coarse powder . . . 4 oz.

Rectified Spirit of Wine . . . 1 pint.

Digest for three days, and then strain.

Used as an emulsifying agent.

UNGUENTUM ACIDI BORACICI.

Boracic Acid Ointment.

Boracic Acid 1 dr.

Oil of Almonds 4 fl. drs.

White Wax 1 dr.

Lard 1 dr.

Mix.

UNGUENTUM ATROPIÆ SULPHATIS.

*Sulphate of Atropia Ointment.*Sulphate of Atropia $\frac{1}{2}$ gr.

Vaseline 1 dr.

Mix.

UNGUENTUM ÆRUGINIS.

Ointment of Subacetate of Copper.

Subacetate of Copper 40 grs.

Prepared Lard 1 oz.

Mix thoroughly.

UNGUENTUM DATURINÆ.

Daturine Ointment.

Daturine	.	.	.	.	.	$\frac{1}{2}$ gr.
Vaseline	.	.	.	.	.	1 dr.

Mix.

UNG. DATURINÆ CUM HYDRARGYRO.

Mercury and Daturine Ointment.

Red Oxide of Mercury	.	.	.	.	.	$\frac{1}{10}$ gr.
Daturine	.	.	.	.	.	$\frac{1}{10}$ gr.
Vaseline	.	.	.	.	.	1 dr.

Mix.

UNGUENTUM DIACHYLI.

Hebra's Diachylon Ointment.

Litharge	.	.	.	.	.	120 grs.
Olive Oil	.	.	.	.	.	1 fl. oz.

Boil with a sufficient quantity of Water, then add


Oil of Lavender	.	.	.	.	.	5 m.
-----------------	---	---	---	---	---	------


UNGUENTUM ESERINÆ.

Eserine Ointment.

Eserine	.	.	.	.	.	$\frac{1}{5}$ gr.
Vaseline	.	.	.	.	.	1 dr.

Mix,





UNGUENTUM HYDRARGYRI NITRATIS
MITIUS.

Milder Nitrate of Mercury Ointment.

Nitrate of Mercury Ointment	.	34	grs.
Lard	.	1	oz.

Mix.

UNGUENTUM HYDRARGYRI OXIDI
FLAVI.

Yellow Oxide of Mercury Ointment.

Yellow Oxide of Mercury	.	2	grs.
Vaseline	.	1	dr.

Mix.

UNGUENTUM IODOFORMI.

Iodoform Ointment.

Iodoform	.	$\frac{1}{2}$	dr.
Simple Ointment	.	1	oz.

Mix.

UNGUENTUM LIQUORIS CARBONIS
DETERGENTIS.

Ointment of Spirit of Coal Tar.

"Liquor Carbonis Detergens"	.	2	fl. drs.
Lard	.	1	oz.

Mix.

UNGUENTUM METALLORUM.

Ointment of Metals.

Ointment of Zinc	} equal parts.
Ointment of Nitrate of Mercury	
Ointment of Acetate of Lead	

Mix.

UNGUENTUM MORPHIÆ ACETATIS.

Acetate of Morphia Ointment.

Acetate of Morphia	1 gr.
Vaseline	1 oz.

Mix.

UNGUENTUM OPII.

Opium Ointment.

Extract of Opium	30 grs.
Prepared Lard	1 oz.
Warm Water	a sufficiency.

Soften the Opium with a little Water, then add the Lard.

UNGUENTUM STAPHYSAGRIÆ.

Ointment of Stavesacre.

Seeds of Stavesacre, crushed	110 grs.
White Wax	55 grs.
Prepared Lard	1 oz.

Melt the Lard and the Wax over a water bath, then add the Stavesacre, and digest while hot for an hour; strain, and stir till cool.

For light Sweats
Luce Oidi gr $\frac{ij}{ij}$ o. n.

To produce vomiting in children

Pulv. Ipecac gr xv. - 3j

Ant. Tart. gr $\frac{1}{2}$. 3/4

Syrup Scilla 3j

Aq. ad. 3ij

3ij ev. 10 min til vomiting

For cleaning sponges
Soda Supt $\mathfrak{Z}\text{ij}$
Acid. Hyd. For $\mathfrak{Z}\text{vi}$.

For cleaning tracheotomy tubes
Bicarb. Soda. gr. xv to the $\mathfrak{Z}\text{i}$
Soak in boiling water, silver tubes.

Cocain Hyd. gr. x
Vaseline Acidi Borici ad $\mathfrak{Z}\text{i}$
in collapsible tube.

Ung Acidi Borici $\mathfrak{Z}\text{i}$ { For burns
Ung vaseline $\mathfrak{Z}\text{i}$

Acidi Tannici gr. xv
Acidi Carbolici gr. x For back
Spt. Rect ad $\mathfrak{Z}\text{i}$, p.a.

UNGUENTUM STYRACIS.

Storax Ointment.

Prepared Storax	145 grs.
Rectified Spirit of Wine	3 fl. drs.
Lard	1 oz.

Mix.

UNGUENTUM ZINCI CUM PLUMBO.

Lead and Zinc Ointment.

Ointment of Zinc	} equal parts.
Ointment of Acetate of Lead	

Mix.

VASELINE.

*A Neutral Hydrocarbon, prepared from
American Petroleum.*

Used in the Ophthalmic Ointments.

DIET TABLE FOR PATIENTS

Full Diet.

Fourteen Ounces of Bread.

One Ounce of Butter.

One Pint of Porter, for Males.
Half-pint of Porter, for
Females. Milk may be sub-
stituted.

Six Ounces of Dressed Meat,
Roast and Boiled, alter-
nately, with half-pound of
Potatoes.

Half-pint of Mutton Broth,
in addition, on days when
Boiled Meat is given.

Half-pound of Baked Rice
Pudding made with Milk,
on Roast Meat days.

Middle Diet.

Twelve Ounces of Bread.

One Ounce of Butter.

Half-pint of Porter for Males
and Females. Milk may be
substituted.

Four Ounces of Dressed Meat,
Roast and Boiled, alter-
nately, with half-pound of
Potatoes.

Half-pint of Mutton Broth,
in addition, on days when
Boiled Meat is given.

Half-pound of Baked Rice
Pudding made with Milk,
on Roast Meat days.

only to be specially ordered

Anemism Diet.

Bread 3 iv
Meat 3 ii
Potatoes 3 ii
Fluid 38

Lemonade (Gav)

Take the juice of a lemon, & add 6 oz of water or soda water. Sugar may be added, as agreeable & a small piece of rind for flavoring.

Take a lemon, peel it carefully, removing all the white part. Cut it into thin slices, remove the pips, add a small slice of thin rind, & pour over it of boiling water, adding sugar as desired. When cold, strain.

Beef tea

Lean Beef, finely minced 1 lb.
Place in a jar & cold water ad of
Jar & allow to stand for 1 hour.
Place jar in saucepan of water & boil gently
for 1 hr, or put jar into warm oven for
same time. Pour the contents of jar on to a
strainer. The B.T. which runs through should
be flavoured & salt. Skin off any fat.
add water to make of

Raw Meat Pulp

Take lean beef or mutton - cut it into
small pieces, & reduce to pulp & pestle
& mortar. Press through a fine sieve, & keep
that on the under surface of sieve, & add
to what has come through, & season the
mixture with a little sugar - Dose for child
of 12 months - 1 tea-spoon.

IN GUY'S HOSPITAL.

Low Diet.

Ten Ounces of Bread.

One Ounce of Butter.

Chop or Fish, with half-pound
of Potatoes.

Beef Tea.

Eggs.

Milk.

Extra Milk and Eggs may
be ordered with Full or Middle
Diet.

Chicken and other delicacies
will be given with the Low
Diet when specially ordered.

Porridge, Gruel, and Barley
Water are given with all Diets,
as required.

Wines and Spirits, if con-
tinued, must be mentioned
each time the Physician or
Surgeon attends.

Tea and Sugar morning and evening with all Diets.

Sister Cornelius.
Nurse Charles.
Nurse Bones.

Emily.
Mrs Martin.

PRINTED BY BALLANTYNE AND HANSON
LONDON AND EDINBURGH



